

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 December 2003 (18.12.2003)

PCT

(10) International Publication Number
WO 03/104180 A1

- (51) International Patent Classification⁷: **C07C 51/347, 57/30**
- (21) International Application Number: **PCT/IN03/00159**
- (22) International Filing Date: **16 April 2003 (16.04.2003)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
427/MAS / 2002 5 June 2002 (05.06.2002) IN
- (71) Applicants (*for all designated States except US*): **NATCO PHARMA LIMITED [IQ/IN]**; Natco House, Road No.2, Banjara Hills, 500033 Hyderabad (IN). **VENKAIAH CHOWDARY NANNAPANENI [IN/IN]**; Natco Pharma Ltd., Natco House, Road No.2, Banjara Hills, 500033 Hyderabad (IN).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): **PULLA REDDY, Muddasani [IN/IN]**; NATCO PHARMA LTD, NATCO HOUSE, ROAD NO.2, BANJARA HILLS, 500033 HYDERABAD, ANDHRA PRADESH (IN).
- (74) Agent: **PODILI KHADGAPATHI**; Natco House, Road No. 2, Banjara Hills, 500033 Hyderabad (IN).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- *with international search report*
 - *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PROCESS FOR THE PREPARATION OF 4-(4-FLUOROBENZOYL) BUTYRIC ACID




(57) Abstract: The present invention provides an improved process for the preparation of 4-(4-Fluorobenzoyl)butyric acid of formula (I), which is prepared on a commercial scale using normal quality fluorobenzene (benzene content 300-700ppm) with the desfluoro analogue impurity at an acceptable level (less than 0.1 % by HPLC). The 4-(4-fluorobenzoyl)butyric acid has the formula (I) is a key raw material for the synthesis of anti-hyperlipoproteinemic drug ezetimibe.

PROCESS FOR THE PREPARATION OF 4-(4-FLUOROBENZOYL) BUTYRIC ACID

5 INTRODUCTION

The present invention relates to an improved process for the preparation of 4-(4-fluorobenzoyl)butyric acid. The 4-(4-fluorobenzoyl)butyric acid has the formula-I given below.



I

10

The 4-(4-fluorobenzoyl)butyric acid has the formula-I is a key raw material for the synthesis of anti-hyperlipoproteinemic drug ezetimibe (US 5767115, Schering). The main criticality in making the compound of formula-I lies in controlling the desfluoro analogue impurity (4-benzoylbutyric acid) at an acceptable level (< 0.05% w/w by HPLC).

15

BACKGROUND OF THE INVENTION

20 In the literature only two procedures are known for the preparation of compound of formula-I. In the first procedure reported by Compennolle (Tetrahedron, 49, 3193, 1993) fluorobenzene is reacted with glutaric anhydride under Friedel-Crafts conditions using aluminium chloride at 0°C to get the compound of formula-I in 78% yield. The reaction is done in methylene chloride medium. In this procedure reaction was done on a 15gr
25 glutaric anhydride scale. The quality of final product with respect to impurities is not addressed in this reference. The main drawbacks in this procedure are:

1. Temperature of the reaction (0°C) is not suitable for scale up operations. At 0°C aluminum chloride-glutaric anhydride complex is not soluble in the medium and a thick mass (difficult to stir by mechanical stirrer) will form. Because of improper mixing quality and yield of the product is low.
 2. Quenching of Friedel-Crafts reaction by adding water or acid to the reaction mass is difficult on a larger scale.
 3. Quality of fluorobenzene used in the process is not mentioned. This is very critical to get an acceptable quality of compound of formula-I with respect to desfluoro analogue impurity.
- 15 In the second route (US patent, 6,207,822) for the synthesis of compound of formula-I, fluorobenzene is used as a solvent-cum-reagent in the Friedel-Crafts acylation with glutaric anhydride. The yield of compound of the formula-I reported in this procedure is 79%. The reaction is done at 5-12°C. The main disadvantages in this process are:
1. It requires high quality (benzene content in fluorobenzene should be less than 100ppm) fluorobenzene to get acceptable quality compound of formula-I.
 2. Availability of high quality fluorobenzene is limited for employing the process on a commercial scale.
 3. Price of high quality fluorobenzene is almost 4 times higher than the normal quality (benzene content is 300-700ppm) fluorobenzene. Normal quality fluorobenzene is abundantly available in the market.

Keeping in view of the above mentioned difficulties in implementing the above routes for making compound of formula-I on a commercial scale, we directed our research work to develop a simple, convenient, and economical process for the preparation of compound of formula-I which can also be utilized on a commercial scale

5

The main objective of the present invention is, therefore, to provide an improved process for the preparation of compound of formula-I as defined above overcoming all the disadvantages present in the hitherto known processes.

- 10 Another objective of the present invention is to provide an improved process for the preparation of compound of formula-I as defined above which is simple and economical.

Another objective of the present invention is to provide an improved process for the preparation of compound of formula-I as defined above which does not have any mixing
15 problem when the process is used on any scale of operation.

20

Another objective of the present invention is to provide an improved process for the preparation of compound of formula-I as defined above which does not have reaction quenching problem when the process is used on any scale of operation.

Still another objective of the present invention is to provide an improved process for the preparation of compound of formula-I as defined above which does not require high quality fluorobenzene.

- 25 Another objective of the present invention is to provide an improved process for the preparation of compound of formula-I as defined above which does not require fluorobenzene as solvent medium to carry out the Friedel-Crafts reaction.

SUMMARY OF INVENTION

In our preliminary studies in the course of the R&D towards development of an improved process for the preparation of compound of formula-I as defined above we found that
5 benzene is more reactive (about 5 times) than fluorobenzene in the above mentioned Friedel-Crafts acylation. This indicates that benzene present in fluorobenzene will react faster than fluorobenzene. Considering this observation we found that

(i) by adjusting the ratio of fluorobenzene with regard to the glutaric anhydride used,


10 (ii) dividing the amount of fluorobenzene used into two parts - one along with glutaric anhydride and another part with the halogenated solvent and

(iii) fixing the quantity of the halogenated solvent used,

a process can be developed for the preparation of compound of the formula-I as defined above which overcomes the drawbacks of the hitherto known processes described above,

15 Further the process developed can also be useful for any commercial scale production of the said compound of the formula-I. The reaction can be conducted at a convenient temperature range of 10 to 25°C.

Accordingly, the present invention provides an improved process for the preparation of
20 compound (4-(4-fluorobenzoyl)butyric acid) of the formula-I



I

which comprises:

25 (a) Preparing a solution of normal quality fluorobenzene, glutaric anhydride and halogenated solvent, the amount of fluorobenzene used being in a molar ratio of 0.5 to 0.7 molar equivalent with regard to the amount of glutaric anhydride used.

- 5 (b) Preparing a mixture of aluminum chloride, normal quality fluorobenzene and halogenated solvent, the amount of fluorobenzene used being in a molar ratio of 0.5 to 0.6 molar equivalent with regard to the amount of glutaric anhydride used and the amount of halogenated solvent used being at least 4-6 times (w/v) with regard to the amount of glutaric anhydride used.
- (c) Adding the solution obtained in step (a) to the mixture obtained in step (b) at a temperature in the range of 10 to 25°C.
- (d) Maintaining the resulting reaction mixture at a temperature in the range of 10 to 25°C for a period in the range of 2 to 4hrs.
- 10 (e) Pouring the reaction mixture into cold dilute hydrochloric acid.
- (f) Distilling the halogenated solvent at atmospheric pressure for its recovery.
- (g) Filtering and washing the residue with the same halogenated solvent used in step (b) above to obtain the compound of the formula-I.
- 15 (h) Purifying the compound of the formula-I by dissolving it in aqueous base and precipitating the product by acidification after giving a carbon treatment to the basic solution.
- (i) Isolating the precipitated compound of formula-I by filtration and if desired
- (j) Recrystallizing the purified acid from a single or mixture of solvents.
- 20 The normal quality of fluorobenzene used in step (a) refers to the impurity level of benzene. The benzene content in fluorobenzene may be between 300-700ppm. The halogenated solvent used in step (b) may be selected from methylene chloride, ethylene dichloride, 1,1,2,2-tetrachloroethane. The base used in step (h) may be selected from sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate,
- 25 sodium hydroxide, potassium hydroxide, and ammonia. The acid used in step (h) may be selected from hydrochloric acid, hydrobromic acid, sulfuric acid, acetic acid, and propionic acid. The solvent used for recrystallization in step (j) may be selected from acetone, methyl ethyl ketone, methyl isobutyl ketone, toluene, acetonitrile, methanol, ethanol, ethyl acetate, hexane or a mixture of these solvents.

The details of the invention are described in Examples given below which are provided to illustrate the invention only and therefore should not be construed to limit the scope of the present invention.

Example 1

Preparation of 4-(4-fluorobenzoyl)butyric acid of formula-I using fluorobenzene (benzene content 300ppm) with methylene chloride as solvent:

10 Into a 3L three-necked RB flask were charged 500ml of methylene chloride, 250gr of aluminum chloride and 45gr of fluorobenzene (benzene content 300ppm) under nitrogen atmosphere. The reaction mixture was cooled to 10°C and a solution of 100gr of glutaric anhydride, 45gr of fluorobenzene (benzene content 300ppm) and 500ml of methylene chloride was added slowly over a period of 3hrs between 10-15°C. The reaction mixture
15 was maintained for another one hour at the same temperature. The reaction mixture was slowly poured onto a mixture of crushed ice (700gr) and conc. HCl (300ml) below 10°C. The reaction mass temperature was allowed to reach 25°C and methylene chloride distilled off from the reaction mixture below 50°C. After cooling the reaction mixture to 20°C, solids were filtered off and washed with 500ml of water.

20

The wet cake thus obtained was suspended in 250-300ml of methylene chloride and filtered. The solid compound was dissolved in 600ml of 4% sodium hydroxide, treated with 10gr of activated charcoal and filtered. The filtrate was acidified with conc. HCl and the precipitated acid was filtered. After washing the wet cake with 500ml of water, it was
25 dissolved in 500ml of acetone. The acetone solution was slowly cooled to 15-20°C and the solid filtered, washed with chilled acetone (50ml) and dried at 50-70°C to get 122gr of white crystalline solid, m.p. 143°C. Purity by HPLC is 99.65%. Desfluoro impurity is less than 0.05%.

Example 2

Preparation of 4-(4-fluorobenzoyl)butyric acid of formula-I using
5 **fluorobenzene (benzene content 500ppm) with methylene chloride as**
solvent:

Into a 3L three-necked RB flask were charged 500ml of methylene chloride, 250gr of
aluminum chloride and 45gr of fluorobenzene (benzene content 500ppm) under nitrogen
10 atmosphere. The reaction mixture was cooled to 10°C and a solution of 100gr of glutaric
anhydride, 45gr of fluorobenzene (benzene content 500ppm) and 500ml of methylene
chloride was added slowly over a period of 3hrs between 10-15°C. The reaction mixture
was maintained for another one hour at the same temperature. The reaction mixture was
slowly poured onto a mixture of crushed ice (700gr) and conc. HCl (300ml) below 10°C.
15 The reaction mass temperature was allowed to reach 25°C and methylene chloride
distilled off from the reaction mixture below 50°C. After cooling the reaction mixture to
20°C, solids were filtered off and washed with 500ml of water.

The wet cake thus obtained was suspended in 250-300ml of methylene chloride and
20 filtered. The solid compound was dissolved in 600ml of 4% sodium hydroxide, treated
with 10gr of activated charcoal and filtered. The filtrate pH was adjusted to 1.0-2.0 with
conc. HCl and the precipitated acid of formula-I was filtered. After washing the wet cake
with 500ml of water, it was dissolved in 500ml of acetone. The acetone solution was
slowly cooled to 15-20°C, maintained for 2h, and the solid filtered, washed with chilled
25 acetone (50ml) and dried at 50-70°C to get 120gr of white crystalline solid of formula-I,
m.p. 143-143.5°C. Purity by HPLC is 99.7%. Desfluoro impurity is less than 0.05%.

Example 3

**Preparation of 4-(4-fluorobenzoyl)butyric acid of formula-I using
5 fluorobenzene (benzene content 700ppm) with methylene chloride as
solvent:**

Into a 3L three-necked RB flask were charged 500ml of methylene chloride, 250gr of
aluminum chloride and 45gr of fluorobenzene (benzene content 700ppm) under nitrogen
10 atmosphere. The reaction mixture was cooled to 10°C and a solution of 100gr of glutaric
anhydride, 45gr of fluorobenzene (benzene content 700ppm) and 500ml of methylene
chloride was added slowly over a period of 3hrs between 10-15°C. The reaction mixture
was maintained for another one hour at the same temperature. The reaction mixture was
slowly poured onto a mixture of crushed ice (700gr) and conc. HCl (300ml) below 10°C.
15 The reaction mass temperature was allowed to reach 25°C and methylene chloride
distilled off from the reaction mixture below 50°C. After cooling the reaction mixture to
20°C, solids were filtered off and washed with 500ml of water.

The wet cake thus obtained was suspended in 250-300ml of methylene chloride and
20 filtered. The solid compound was dissolved in 600ml of 4% sodium hydroxide, treated
with 10gr of activated charcoal and filtered. The filtrate pH was adjusted to 1.0-2.0 with
conc. HCl and the precipitated acid of formula-I was filtered. After washing the wet cake
with 500ml of water, it was dissolved in 500ml of acetone. The acetone solution was
slowly cooled to 15-20°C, maintained for 2h, and the solid filtered, washed with chilled
25 acetone (50ml) and dried at 50-70°C to get 123gr of white crystalline solid of formula-I,
m.p. 143°C. Purity by HPLC is 99.6%. Desfluoro impurity is less than 0.05%.

Example 4

**Preparation of 4-(4-fluorobenzoyl)butyric acid of formula-I using
5 fluorobenzene (benzene content 500ppm) with ethylene dichloride as
solvent:**

Into a 3L three-necked RB flask was charged ethylene dichloride (500ml), aluminum
chloride (250gr) and fluorobenzene (45gr) under nitrogen atmosphere. The reaction
mixture was cooled to 10°C and a solution of glutaric anhydride (100gr), fluorobenzene
10 (45gr) and ethylene dichloride (500ml) was added slowly over a period of 3hrs between
10-15°C. After maintaining for one hour at 15-18°C the reaction mixture was slowly
poured onto a mixture of crushed ice (700gr) and conc. HCl (300ml) below 10°C. The
reaction mass temperature was raised to reach 25°C and distilled off ethylene dichloride
from the reaction mixture below 100°C. After cooling the reaction mixture to 20°C, crude
15 solid was filtered off and washed with 500ml of water.


The wet cake thus obtained was suspended in 300ml of ethylene dichloride and filtered.
The solid compound was dissolved in 600ml of 4% sodium hydroxide, treated with 15gr
of activated charcoal and filtered. The filtrate was acidified to pH 1.5-2.0 with conc. HCl
20 and the precipitated acid was filtered. After washing the wet cake with 500ml of water, it
was dried at 60-70°C to get 130gr of white solid, m.p. 140-142°C. This solid was
dissolved in 500ml of acetone. The acetone solution was slowly cooled to 15-20°C and
the solid filtered, washed with chilled acetone (50ml) and dried at 50-70°C to get 120gr
of white crystalline solid, m.p. 143°C. Purity by HPLC is 99.65%. Desfluoro impurity is
25 less than 0.05%.

Advantages of the present invention:

1. (a) By adjusting the ratio of fluorobenzene with regard to the glutaric anhydride used
(b) dividing the amount of fluorobenzene used into two parts - one along with glutaric
5 anhydride and another part with the halogenated solvent and (c) conducting the
Friedel-Crafts acylation at 10-20°C, the process can be made applicable to any scale
of operation.
2. By using halogenated solvent in the reaction, quality of required fluorobenzene could
10 be relaxed to normal quality to make the process economical.
3. High quality compound of formula-I could be produced using normal quality
fluorobenzene.

We Claim:

1. An improved process for the preparation of compound of formula-I

**I**

which comprises:

- (a) Preparing a solution of normal quality fluorobenzene, glutaric anhydride and halogenated solvent, the amount of fluorobenzene used being in a molar ratio of 0.5 to 0.7 molar equivalent with regard to the amount of glutaric anhydride used.
- (b) Preparing a mixture of aluminum chloride, normal quality fluorobenzene and halogenated solvent, the amount of fluorobenzene used being in a molar ratio of 0.5 to 0.6 molar equivalent with regard to the amount of glutaric anhydride used and the amount of halogenated solvent used being at least 4-6 times (w/v) with regard to the amount of glutaric anhydride used.
- (c) Adding the solution obtained in step (a) to the mixture obtained in step (b) at a temperature in the range of 10 to 25°C.
- (d) Maintaining the reaction mixture at the temperature in the range of 10 to 25°C for a period in the range of 2 to 4hrs.
- (e) Pouring the reaction mixture into cold dilute hydrochloric acid.
- (f) Distilling the halogenated solvent at atmospheric pressure for its recovery.
- (g) Filtering and washing the residue with the same halogenated solvent used in step (b) above to obtain the compound of the formula-I.
- (h) Purifying the compound of the formula-I by dissolving it in aqueous base and precipitating the product by acidification after giving a carbon treatment to the basic solution.

- (i) Isolating the precipitated compound of formula-I by filtration and if desired
- (j) Recrystallizing the purified acid from a single or mixture of solvents.

2. An improved process for the preparation of compound of formula-I as claimed in
5 claim 1(i) wherein the normal quality fluorobenzene used in the process has a
benzene content of 300-700ppm, preferably between 300-500ppm.
3. An improved process for the preparation of compound of formula-I as claimed in
10 claims 1 & 2 wherein the halogenated solvent used in the reaction is methylene
chloride, ethylene dichloride, 1,1,2,2-tetrachloroethylene, preferably methylene
chloride or ethylene dichloride.
4. An improved process for the preparation of compound of formula-I as claimed in
15 claims 1 to 3 wherein the quantity of solvent used is 6 to 10 times (w/v) on glutaric
anhydride, preferably 8 to 10 times.
5. An improved process for the preparation of compound of formula-I as claimed in
claims 1 to 4 wherein the reaction temperature is between 10-25°C, preferably
between 12-18°C.
- 20 6. An improved process for the preparation of compound of formula-I as claimed in
claims 1 to 5 wherein the base used to dissolve the crude acid is ammonia, sodium
carbonate, sodium bicarbonate, sodium hydroxide, potassium carbonate, potassium
bicarbonate, potassium hydroxide, ammonia, preferably sodium hydroxide or
25 ammonia.
7. An improved process for the preparation of compound of formula-I as claimed in
claims 1 to 6 wherein the acid used to neutralize the base is hydrochloric acid,

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IN 03/00159

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07C51/347 C07C57/30

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, PAJ, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 38305 A (BARTH MARTINE ;DODEY PIERRE (FR); PAQUET JEAN LUC (FR); PRUNEAU DI) 31 May 2001 (2001-05-31) *preparation IV*	1-7
A	BAENS, NICOLE P. ET AL: "Synthesis of 2,5-substituted piperidines: transposition of 1,4-substitution pattern for the analgesic drug R6582" TETRAHEDRON (1993), 49(15), 3193-202 , XP002022790 page 3197	1-7
A	WO 00 34240 A (SCHERING CORP) 15 June 2000 (2000-06-15) *page 11*	1-7

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

15 September 2003

Date of mailing of the international search report

10/10/2003

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lorenzo Varela, M.J.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IN 03/00159

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>US 6 207 822 B1 (CHIU JOHN S ET AL) 27 March 2001 (2001-03-27) *columns 12 and 13*</p>	1-7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IN 03/00159

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0138305	A	31-05-2001	FR 2801585 A1	01-06-2001
			AU 2177601 A	04-06-2001
			BR 0015695 A	23-07-2002
			CA 2392225 A1	31-05-2001
			CN 1391556 T	15-01-2003
			CZ 20021810 A3	14-08-2002
			EP 1232144 A2	21-08-2002
			WO 0138305 A2	31-05-2001
			HU 0203603 A2	28-02-2003
			JP 2003514894 T	22-04-2003
			NO 20022460 A	24-05-2002
			SK 6892002 A3	08-10-2002
			US 6605633 B1	12-08-2003
			WO 0034240	A
CA 2353981 A1	15-06-2000			
CN 1329592 T	02-01-2002			
EP 1137634 A1	04-10-2001			
JP 2002531546 T	24-09-2002			
WO 0034240 A1	15-06-2000			
ZA 200104004 A	16-08-2002			
US 6207822	B1	27-03-2001	NONE	